Global Search for New Physics in 2 fb^{-1} at CDF

Si Xie

FNAL Users' Meeting
June 5 , 2008

Standard Model Works Very Well



We expect something new !



NOT YET THOUGHT OF
NOT YET THOUGHT OF
NOT THOUGHT
TC-TC
composite Higgs
hypercolor
supercolor
technI-GIM
extended TC
pseudo NGB
6th gen
5th gen
4th gen
S.T.U
vector-like family
lepto quark
heavy Majorana
fractionally charged
sterile ν
milli-charged
shadow matter
triplet Higgs
general 2HDM
Type II
spontaneous CP
superweak
milli-weak
weinberg's 3HD
Majoron
axion
familon
NGB
majoron
extra dim
quintessence
composite W, Z
contact
string
MSSM
NMSSM


bb bb
jj bb
jj jj
γ bb
γ jj
γ γ
τ bb
τ jj
τ γ
τ τ
μ bb
μ jj
μ γ
μ τ
μ μ
e bb
e jj
e γ
e τ
e μ
e e
bb
jj
γ
τ
μ
e
Look everywhere
LQ
ρ_T
Z'
W'
e μ τ $\not{p}_T$ γ jj bb ee eμ eτ e$\not{p}_T$ eγ ejj ebb μμ μτ μ$\not{p}_T$ μγ μjj μbb ττ τ$\not{p}_T$ τγ τjj τbb $\not{p}_T\gamma$ $\not{p}_T$jj $\not{p}_T$bb γγ γjj γbb

Strategy Overview

- Analyze as many data events as computing ability allows
- Construct global Standard Model background prediction
- Search for discrepancies between data and standard model prediction
- Focus attention on outliers (“5σ effects”) !!!

For reference the method is described in detail here: arXiv:0712.1311 accepted by Phys Rev D

Identify Objects and Select Events

- Identify Physics Objects

$p_T > 17\text{GeV}$



- Select Events of Interest
  - Select events containing high-p_T objects, diobjects, multi-objects
  - Total Selection of ~4 million events

Full Tevatron Standard Model Monte Carlo Set

Dataset	Process	Weights	*	Number	=	Total weight
pyth_jj_000	Pythia jj 0<pT<10	1100		2		2113.78
pyth_jj_010	Pythia jj 10<pT<18	500		57		28500
pyth_pj_008	Pythia j gamma 8<pT<12	87		5		434.31
mrenna_mu+mu-	MadEvent Z(-> mu mu)	30		219		6474.94
pyth_jj_090	Pythia jj 90<pT<120	22		2035		45680.6
pyth_pj_012	Pythia j gamma 12<pT<22	21		1974		42110.7
pyth_jj_018	Pythia jj 18<pT<40	19		23898		450480
mad_vtvt-j	MadEvent Z(->vv) j	16		2		31.86
mad_vevе-j	MadEvent Z(->vv) j	16		2		31.69
alpgen_eve	Alpgen W(->e v)	12		5823		69289.9
mrenna_e+e-	MadEvent Z(->ee)	10		5974		60159.9
alpgen_muve	Alpgen W(-> mu v)	9.9		4483		44213.6
pyth_jj_120	Pythia jj 120<pT<150	8.3		3282		27170.8
pyth_jj_060	Pythia jj 60<pT<90	6.7		25299		170363
mrenna_mu+mu-j	MadEvent Z(-> mu mu) j	6.6		3211		21126
pyth_jj_040	Pythia jj 40<pT<60	5		88450		438739
pyth_bj_010	Pythia bj 10<pT<18	3.6		167		604.26
pyth_jj_200	Pythia jj 200<pT<300	3.4		72998		249296
mad_vevе-a_f	MadEvent Z(->vv) gamma	3.4		13		44.23
ut0sw0	Alpgen W(-> tau v)+jets	3.2		649		2063.06
pyth_pj_022	Pythia j gamma 22<pT<45	3		31308		94944
pyth_jj_150	Pythia jj 150<pT<200	2.7		59222		162273
we0sfe	Pythia W(->e v)	2.4		381176		920761
cosmic_j_lo	Cosmic (jet20)	2.3		122		276.85
cosmic_ph	Cosmic (photon_25_iso)	1.8		2700		4892.78
pyth_pj_080	Pythia j gamma 80<pT	1.5		18464		28033.3
mrenna_e+e-j	MadEvent Z(->ee) j	1.4		28137		40761
pyth_pj_045	Pythia j gamma 45<pT<80	1.4		83370		117889
mrenna_mu+mu-jj	MadEvent Z(-> mu mu) jj	1.3		4150		5503.82
pyth_bj_018	Pythia bj 18<pT<40	1.1		16076		18233.3
mad_e+e-	MadEvent Z(->ee)	1		522		542.22
stelzer_l+l-j	stelzer_l+l-j	0.92		665		611.86
mrenna_e+e-jj	MadEvent Z(->ee) jj	0.91		11292		10317.9
mad_mu+mu-	MadEvent Z(-> mu mu)	0.88		83		73.28
pyth_bj_060	Pythia bj 60<pT<90	0.87		10711		9307.8
mad_vtvt-a_f	MadEvent Z(->vv) gamma	0.85		38		32.2
pyth_bj_090	Pythia bj 90<pT<120	0.83		2385		1985.66
mad_vtvt-j_f	MadEvent Z(->vv) j	0.71		7		4.94
stelzer_Waj	MadEvent W(->l v)j gamma	0.68		1644		1125.1
pyth_bj_120	Pythia bj 120<pT<150	0.67		2854		1904.7
mad_aaj	MadEvent j gamma gamma	0.51		563		287.44
we0s8m	Pythia W(-> mu v)	0.49		1.2908e+06		630864
pyth_bj_150	Pythia bj 150<pT<200	0.44		28229		12531.9
mrenna_mu+mu-jjj	MadEvent Z(-> mu mu) jjj	0.44		3448		1530.61
mad_e+e-j	MadEvent Z(->ee) j	0.39		733		285.76
alpgen_evej	Alpgen W(->e v) j	0.35		398712		140567
we0sat	Pythia W(-> tau v)	0.35		49498		17125.5
mad_mu+mu-j	MadEvent Z(-> mu mu) j	0.34		495		166.31
mad_mu+mu-jj	MadEvent Z(-> mu mu) jj	0.32		1682		531.82
ze1s0m	Pythia Z(-> mu mu)	0.3		371008		110522
alpgen_muvej	Alpgen W(-> mu v) j	0.3		281049		83604.3

ut0s2w	Alpgen W(-> tau v)+jets	0.29	5220	1534.87
mad_vtvt-a	MadEvent Z(->vv) gamma	0.27	138	37.48
mad_vevе-a	MadEvent Z(->vv) gamma	0.27	139	37.39
we0s9t	Pythia W(-> tau v)	0.26	66004	17092.3
ut0swi	Alpgen W(-> tau v)+jets	0.24	27810	6632.37
pyth_pp	Pythia gamma gamma	0.23	25786	5807.24
ze1s6d	Pythia Z(->ee)	0.22	484911	106271
mad_e+e-b-b	MadEvent Z(->ee) bb	0.22	1031	224.6
re0s28	Baur W(->ev) gamma	0.21	22076	4701.49
alpgen_evejj	Alpgen W(->e v) jj	0.21	175607	37356.5
alpgen_muvmjj	Alpgen W(-> mu v) jj	0.2	112548	22156.7
ztopcz	Pythia ZZ	0.19	583	109.58
stelzer_Zaj	stelzer_Zaj	0.18	1585	286.94
mad_aajj	MadEvent jj gamma gamma	0.18	7872	1415.27
mad_mu+mu-b-b	MadEvent Z(-> mu mu) bb	0.18	619	108.52
mad_e+e-jj	MadEvent Z(->ee) jj	0.17	773	133.82
re0s29	Baur W(-> mu v) gamma	0.17	19999	3461.88
re0s1a	Baur W(-> tau v) gamma	0.17	2837	488.24
mad_vevе-j_f	MadEvent Z(->vv) j	0.16	14	2.21
pyth_jj_300	Pythia jj 300<pT<400	0.14	103806	14875.4
mad_aaa_f	MadEvent gamma gamma gamma	0.14	55	7.59
cosmic_j_hi	Cosmic (jet100)	0.12	36667	4484.23
pyth_bj_040	Pythia bj 40<pT<60	0.12	161606	18764.2
mrenna_e+e-jjj	MadEvent Z(->ee) jjj	0.11	23968	2661.32
ze0s8t	Pythia Z(-> tau tau)	0.092	16278	1496.71
pyth_bj_200	Pythia bj 200<pT<300	0.081	252367	20555.5
hewk08	MadEvent Z(->ee) gamma	0.081	70511	5713.41
mad_aaa	MadEvent gamma gamma gamma	0.079	72	5.69
zz0sem	Pythia Z(-> mu mu) (m_Z<20)	0.075	30	2.26
wenubb0p	Alpgen W(->e v) bb	0.075	41332	3096.21
wmnubb0p	Alpgen W(-> mu v) bb	0.075	25998	1946.94
zz0see	Pythia Z(->ee) (m_Z<20)	0.074	79	5.85
overlay	Overlaid events	0.073	11443	837.38
wenubbip	Alpgen W(->e v) bb j	0.072	14076	1018.56
wmnubbip	Alpgen W(-> mu v) bb j	0.072	8420	608.96
hewk04	MadEvent Z(-> mu mu) gamma	0.072	2034	145.66
pyth_jj_400	Pythia jj 400<pT	0.068	13106	890.33
alpgen_evejjj	Alpgen W(->e v) jjj	0.068	92568	6259.88
alpgen_muvmjjj	Alpgen W(-> mu v) jjj	0.066	55644	3689.5
ttop0z	Herwig ttbar	0.065	30649	1982.71
ze0sat	Pythia Z(-> tau tau)	0.063	23833	1512.59
ut0s3w	Alpgen W(-> tau v)+jets	0.063	4470	282.34
wmnubb2p	Alpgen W(-> mu v) bb jj	0.064	3508	188.94
wenubb2p	Alpgen W(->e v) bb jj	0.064	6044	323.72
we0scd	Pythia WZ	0.063	2910	154.95
we0sgd	Pythia WW	0.048	2563	122.77
we0sbd	Pythia WW	0.048	2843	136.03
alpgen_evejjjj	Alpgen W(->e v) jjjj	0.027	41589	1118.82
alpgen_muvmjjjj	Alpgen W(-> mu v) jjjj	0.024	26964	650.93
ut0s4w	Alpgen W(-> tau v)+jets	0.023	2488	57.06
Total:				4.37683e+06

Simulate detector response



Correction Model

- Attempts to accurately reflect the limit to our systematic understanding of the detector and the standard model
- Correction factors include: integrated luminosity, k-factors, trigger efficiencies, reconstruction efficiencies, fake rates
- Values are obtained by a global fit of data to background yielding a set of values maximizing global agreement

Some typical distributions that constrain correction factors









Partition Events into **exclusive final states**



Global Comparison

399 Final State Populations

CDF Run II Preliminary ($2.0\ \text{fb}^{-1}$)
The calculation of σ accounts for the trials factor

Final State	Data	Background
$be^{\pm}\not{p}$	690	817.7 ± 9.2
$\gamma\tau^{\pm}$	1371	1217.6 ± 13.3
$\mu^{\pm}\tau^{\pm}$	63	35.2 ± 2.8
b2j$\not{p}$ high-Σp_T	255	327.2 ± 8.9
2j$\tau^{\pm}$ low-Σp_T	574	670.3 ± 8.6
3j$\tau^{\pm}$ low-Σp_T	148	199.8 ± 5.2
$e^{\pm}\not{p}\tau^{\pm}$	36	17.2 ± 1.7
2j$\tau^{\pm}\tau^{\mp}$	33	62.1 ± 4.3
$e^{\pm}$j	741710	$764832 \pm 6447.$
j2$\tau^{\pm}$	105	150.8 ± 6.3
$e^{\pm}$2j	256946	$249148 \pm 2201.$
2bj low-Σp_T	279	352.5 ± 11.9
j$\tau^{\pm}$ low-Σp_T	1385	1525.8 ± 15
2b2j low-Σp_T	108	153.5 ± 6.8
b$\mu^{\pm}\not{p}$	528	613.5 ± 8.7
$\mu^{\pm}\gamma\not{p}$	523	611 ± 12.1
2bγ	108	70.5 ± 7.9
8j	14	13.1 ± 4.4
7j	103	97.8 ± 12.2
6j	653	659.7 ± 37.3
5j	3157	3178.7 ± 67.1
4j high-Σp_T	88546	89096.6 ± 935.2
4j low-Σp_T	14872	14809.6 ± 186.3
4j2γ	46	46.4 ± 3.9
4j$\tau^{\pm}$ high-Σp_T	29	26.6 ± 1.7
4j$\tau^{\pm}$ low-Σp_T	43	63.1 ± 3.3
4j$\not{p}$ high-Σp_T	1064	1012 ± 62.9
4j$\gamma\tau^{\pm}$	19	10.8 ± 2
4j$\gamma\not{p}$	62	104.2 ± 22.4
4jγ	7962	8271.2 ± 245.1
4j$\mu^{\pm}\not{p}$	574	590.5 ± 13.6
4j$\mu^{\pm}\mu^{\mp}$	38	48.4 ± 6.2
4j$\mu^{\pm}$	1363	1350.1 ± 37.7
3j high-Σp_T	159926	$159143 \pm 1061.$
3j low-Σp_T	62681	64213.1 ± 496
3j2γ	151	177.5 ± 7.1
3j$\tau^{\pm}$ high-Σp_T	68	76.9 ± 3
3j$\not{p}$ high-Σp_T	1706	1899.4 ± 77.6
3j$\not{p}$ low-Σp_T	42	36.2 ± 5.7
3j$\gamma\tau^{\pm}$	39	37.8 ± 3.6
3j$\gamma\not{p}$	204	249.8 ± 24.4
3jγ	24639	24899.4 ± 372.4
3j$\mu^{\pm}\not{p}$	2884	2971.5 ± 52.1
3j$\mu^{\pm}\gamma\not{p}$	10	3.6 ± 1.9
3j$\mu^{\pm}\gamma$	15	7.9 ± 2.9
3j$\mu^{\pm}\mu^{\mp}$	175	177.8 ± 16.2
3j$\mu^{\pm}$	5032	4989.5 ± 108.9
3b2j	23	28.9 ± 4.7
3bj	82	82.6 ± 5.7
3b	67	85.6 ± 7.7
2$\tau^{\pm}$	498	512.7 ± 14.2
2$\gamma\not{p}$	128	107.2 ± 6.9
2γ	5548	5562.8 ± 40.5
2j high-Σp_T	190773	190842 ± 781.2
2j low-Σp_T	165984	162530 ± 1581
2j2$\tau^{\pm}$	22	40.6 ± 3.2
2j2$\gamma\not{p}$	11	8 ± 2.4
2j2γ	580	581 ± 13.7
2j$\tau^{\pm}$ high-Σp_T	96	114.6 ± 3.3

Final State	Data	Background
2j$\not{p}$ high-Σp_T	87	80.9 ± 6.8
2j$\not{p}$ low-Σp_T	114	79.5 ± 100.8
2j$\not{p}\tau^{\pm}$	18	13.2 ± 2.2
2j$\gamma\tau^{\pm}$	142	144.6 ± 5.7
2j$\gamma\not{p}$	908	980.3 ± 63.7
2jγ	71364	73021.4 ± 595.9
2j$\mu^{\pm}\tau^{\mp}$	16	19.3 ± 2.2
2j$\mu^{\pm}\not{p}$	17927	18340.6 ± 201.9
2j$\mu^{\pm}\gamma\not{p}$	31	27.7 ± 7.7
2j$\mu^{\pm}\gamma$	57	58.2 ± 13
2j$\mu^{\pm}\mu^{\mp}\not{p}$	11	7.8 ± 2.7
2j$\mu^{\pm}\mu^{\mp}$	956	924.9 ± 61.2
2j$\mu^{\pm}$	22461	23111.4 ± 366.6
2e$^{\pm}$j	14	13.8 ± 2.3
2e$^{\pm}e^{\mp}$	20	17.5 ± 1.7
2e$^{\pm}$	32	49.2 ± 3.4
2b high-Σp_T	666	689 ± 9.4
2b low-Σp_T	323	313.2 ± 10.3
2b3j low-Σp_T	53	57.4 ± 6.5
2b2j high-Σp_T	718	803.3 ± 12.7
2b2j$\not{p}$ high-Σp_T	15	21.8 ± 2.8
2b2jγ	32	39.7 ± 6.2
2b2j$\mu^{\pm}\not{p}$	14	17.3 ± 1.9
2b2j$\mu^{\pm}$	22	21.8 ± 2
2b$\mu^{\pm}\not{p}$	11	14.4 ± 2.1
2bj high-Σp_T	891	967.1 ± 13.2
2bj$\not{p}$ high-Σp_T	25	31.3 ± 3.1
2bjγ	71	54.5 ± 7.1
2bj$\mu^{\pm}\not{p}$	12	10.7 ± 1.9
2be$^{\pm}$2j$\not{p}$	30	27.3 ± 2.2
2be$^{\pm}$2j	72	66.5 ± 2.9
2be$^{\pm}\not{p}$	22	19.1 ± 2.2
2be$^{\pm}$j$\not{p}$	19	19.4 ± 2.2
2be$^{\pm}$j	63	63 ± 3.4
2be$^{\pm}$	96	92.1 ± 4.1
$\tau^{\pm}\tau^{\mp}$	856	872.5 ± 19
$\gamma\not{p}$	3793	3770.7 ± 127.3
$\mu^{\pm}\tau^{\mp}$	381	440.9 ± 7.3
$\mu^{\pm}\not{p}\tau^{\mp}$	60	75.7 ± 3.4
$\mu^{\pm}\not{p}\tau^{\pm}$	15	12 ± 2
$\mu^{\pm}\not{p}$	734290	734296 ± 4897.8
$\mu^{\pm}\gamma$	475	469.8 ± 12.5
$\mu^{\pm}\mu^{\mp}\not{p}$	169	198.5 ± 8.2
$\mu^{\pm}\mu^{\mp}\gamma$	83	60 ± 3.1
$\mu^{\pm}\mu^{\mp}$	25283	25178.5 ± 86.5
j2$\gamma\not{p}$	36	30.4 ± 4.2
j2γ	1822	1813.2 ± 27.4
j$\tau^{\pm}$ high-Σp_T	52	56.2 ± 2.5
j$\tau^{\pm}\tau^{\mp}$	203	252.2 ± 8.7
j$\not{p}$ high-Σp_T	4432	4431.7 ± 45.2
j$\gamma\tau^{\pm}$	526	476 ± 9.3
j$\gamma\not{p}$	1882	1791.9 ± 72.3
jγ	103319	102124 ± 570.6
j$\mu^{\pm}\tau^{\mp}$	71	98 ± 3.9
j$\mu^{\pm}\tau^{\pm}$	15	12 ± 2
j$\mu^{\pm}\not{p}\tau^{\mp}$	26	30.8 ± 2.6
j$\mu^{\pm}\not{p}$	109081	108323 ± 707.7
j$\mu^{\pm}\gamma\not{p}$	171	171.1 ± 31
j$\mu^{\pm}\gamma$	152	190 ± 39.3

Final State	Data	Background
j$\mu^{\pm}\mu^{\mp}\not{p}$	32	32.2 ± 10.9
j$\mu^{\pm}\mu^{\mp}\gamma$	14	11.5 ± 2.6
j$\mu^{\pm}\mu^{\mp}$	4852	4271.2 ± 185.4
j$\mu^{\pm}$	77689	76987.5 ± 930.2
$e^{\pm}$4j$\not{p}$	903	830.6 ± 13.2
$e^{\pm}$4jγ	25	29.2 ± 3.6
$e^{\pm}$4j	15750	16740.4 ± 390.5
$e^{\pm}$3j$\tau^{\mp}$	15	21.1 ± 2.2
$e^{\pm}$3j$\not{p}$	4054	4077.2 ± 63.6
$e^{\pm}$3jγ	108	79.3 ± 5
$e^{\pm}$3j	60725	60409.3 ± 723.3
$e^{\pm}$2γ	41	34.2 ± 2.6
$e^{\pm}$2j$\tau^{\pm}$	37	47.2 ± 2.2
$e^{\pm}$2j$\tau^{\mp}$	109	95.9 ± 6.8
$e^{\pm}$2j$\not{p}$	25725	25403.1 ± 209.4
$e^{\pm}$2j$\gamma\not{p}$	30	31.8 ± 4.8
$e^{\pm}$2jγ	398	342.8 ± 15.7
$e^{\pm}$2j$\mu^{\mp}\not{p}$	22	14.8 ± 1.9
$e^{\pm}$2j$\mu^{\mp}$	23	15.8 ± 2
$e^{\pm}\tau^{\pm}$	437	387 ± 5.3
$e^{\pm}\tau^{\mp}$	1333	1266 ± 12.3
$e^{\pm}\not{p}\tau^{\mp}$	109	106.1 ± 2.7
$e^{\pm}\not{p}$	960826	$956579 \pm 3077.$
$e^{\pm}\gamma\not{p}$	497	496.8 ± 10.3
$e^{\pm}\gamma$	3578	3589.9 ± 24.1
$e^{\pm}\mu^{\pm}\not{p}$	31	29.9 ± 1.6
$e^{\pm}\mu^{\mp}\not{p}$	109	99.4 ± 2.4
$e^{\pm}\mu^{\pm}$	45	28.5 ± 1.8
$e^{\pm}\mu^{\mp}$	350	313 ± 5.4
$e^{\pm}$j2γ	13	16.1 ± 3.9
$e^{\pm}$j$\tau^{\mp}$	386	418 ± 18.9
$e^{\pm}$j$\tau^{\pm}$	160	162.8 ± 3.5
$e^{\pm}$j$\not{p}\tau^{\mp}$	48	44.6 ± 3.3
$e^{\pm}$j$\not{p}\tau^{\pm}$	11	8.3 ± 1.5
$e^{\pm}$j$\not{p}$	121431	121023 ± 747.6
$e^{\pm}$j$\gamma\not{p}$	159	192.6 ± 10.9
$e^{\pm}$jγ	1389	1368.9 ± 38.9
$e^{\pm}$j$\mu^{\mp}\not{p}$	42	33 ± 2.9
$e^{\pm}$j$\mu^{\pm}\not{p}$	16	9.2 ± 1.9
$e^{\pm}$j$\mu^{\mp}$	62	63.8 ± 3.2
$e^{\pm}$j$\mu^{\pm}$	13	8.2 ± 2
$e^{\pm}e^{\mp}$4j	148	159.1 ± 7
$e^{\pm}e^{\mp}$3j	717	743.6 ± 24.4
$e^{\pm}e^{\mp}$2j$\not{p}$	32	41.4 ± 5.6
$e^{\pm}e^{\mp}$2jγ	10	11.4 ± 2.9
$e^{\pm}e^{\mp}$2j	3638	3566.8 ± 72
$e^{\pm}e^{\mp}\tau^{\pm}$	18	16.1 ± 1.7
$e^{\pm}e^{\mp}\not{p}$	822	831.8 ± 13.6
$e^{\pm}e^{\mp}\gamma$	191	221.9 ± 5.1
$e^{\pm}e^{\mp}$j$\not{p}$	155	170.8 ± 12.4
$e^{\pm}e^{\mp}$jγ	48	45 ± 3.9
$e^{\pm}e^{\mp}$j	17903	18258.2 ± 204.4
$e^{\pm}e^{\mp}$	98901	99086.9 ± 147.8
b6j	51	42.3 ± 3.8
b5j	237	192.5 ± 7.1
b4j high-Σp_T	26	23.4 ± 2.6
b4j low-Σp_T	836	821.7 ± 15.9
b3j high-Σp_T	12081	12071 ± 84.1
b3j low-Σp_T	2974	2873 ± 31

Global Comparison

19650 Kinematic Distributions









Global Comparison

Bump Hunter: 5036 Mass variables



- Search for resonant production of new particles
- Look for narrow resonances in mass distributions.
- (width = detector resolution)
- Require quality criteria to eliminate erroneous bumps

Global Comparison

Sleuth Final States : 87

Sleuth variable: $$\sum p_T \equiv \sum_i |\vec{p}_i| + \left|\overrightarrow{\text{uncl}}\right| + \left|\vec{\not{p}}\right|,$$



- Scan the ΣP_T spectrum
- Look for semi-infinite region with the most significant excess of data
- Excesses at Large ΣP_T are expected by a wide spectrum of new physics scenarios.

Overview Schematic




Results

Final State Populations

List of Top Population Discrepancies

CDF Run II Preliminary ($2.0\ \text{fb}^{-1}$)
The calculation of σ accounts for the trials factor

Final State	Data	Background	σ
$be^{\pm}\not{p}$	690	817.7 ± 9.2	-2.7
$\gamma\tau^{\pm}$	1371	1217.6 ± 13.3	$+2.2$
$\mu^{\pm}\tau^{\pm}$	63	35.2 ± 2.8	$+1.7$
b2j$\not{p}$ high-Σp_T	255	327.2 ± 8.9	-1.7
2j$\tau^{\pm}$ low-Σp_T	574	670.3 ± 8.6	-1.5
3j$\tau^{\pm}$ low-Σp_T	148	199.8 ± 5.2	-1.4
$e^{\pm}\not{p}\tau^{\pm}$	36	17.2 ± 1.7	$+1.4$
2j$\tau^{\pm}\tau^{\mp}$	33	62.1 ± 4.3	-1.3
$e^{\pm}$j	741710	764832 ± 6447.2	-1.3
j2$\tau^{\pm}$	105	150.8 ± 6.3	-1.2
$e^{\pm}$2j	256946	249148 ± 2201.5	$+1.2$
2bj low-Σp_T	279	352.5 ± 11.9	-1.1
j$\tau^{\pm}$ low-Σp_T	1385	1525.8 ± 15	-1.1
2b2j low-Σp_T	108	153.5 ± 6.8	-1
b$\mu^{\pm}\not{p}$	528	613.5 ± 8.7	-0.9
$\mu^{\pm}\gamma\not{p}$	523	611 ± 12.1	-0.8
2bγ	108	70.5 ± 7.9	$+0.1$
8j	14	13.1 ± 4.4	0
7j	103	97.8 ± 12.2	0
6j	653	659.7 ± 37.3	0
5j	3157	3178.7 ± 67.1	0



No Significant Discrepancy in Populations!

Kinematic Shapes





This analysis brought this issue ("3 jet effect") to attention and is currently being investigated by experimental and theoretical colleagues.

This is a major limiting factor in our ability to resolve mass resonances in multijet final states.

3 jet effect







Conclusion: No discrepancies to motivate a new physics claim

Summary of Mass Bumps



The outlier





4.1 sigma bump after trials factor



No new physics interpretation. "3-jet" effect again!!!

Sleuth Results

$e^{\pm}\mu^{\pm}$





$e^{\pm}\mu^{\pm} \not{E}_T$ jj

$e^{\pm}\mu^{\pm} \not{E}_T$





$e^+e^-\mu^{\pm} \not{E}_T$

$\mu^+\mu^- e^{\pm} \not{E}_T$

Sleuth Results Summary

CDF Run II Preliminary ($2.0\ \mathrm{fb}^{-1}$)

SLEUTH Final State	$\mathcal{P}$
$\ell^+\ell'^+$	0.00055
$\ell^+\ell'^+ \not{p} jj$	0.0021
$\ell^+\ell'^+ \not{p}$	0.0042
$\ell^+\ell^-\ell' \not{p}$	0.0047
$\ell^+\tau^+ \not{p}$	0.0065

The most discrepant Sleuth final state has a probability of 8% after accounting for trials factor

Conclusion

- Completed the global search for new physics in 2fb^{-1} at CDF
- This search represents one of the most encompassing tests of the Standard Model at the energy frontier.
- This analysis finds...
  - No significant final state population discrepancies.
  - No shape discrepancies motivating new physics
  - No bumps motivating new physics
  - No statistically significant Sleuth Σp_T excess.
- The search continues...

BACKUP

A few example distributions

From
Most significant deficit



From
Most significant excess



Sensitivity: Top Discovery?

Remove top quark from Standard Model, refit correction factors, search!



Easily finding top in 1 fb^{-1}

Approximate Luminosity Needed ~ O (Run1 Discovery)

Sensitivity: Z' mass bump ?



Z'_{250} → charged leptons
5σ discovery if σ×BR ≈ 0.325 pb.

Global Fit

$$\chi^2(\vec{s}) = \left(\sum_{k \in \text{bins}} \chi^2_k(\vec{s}) \right) + \chi^2_{\text{constraints}}(\vec{s})$$

$$\chi^2_k(\vec{s}) = \frac{(\text{Data}[k] - \text{SM}[k])^2}{\delta\text{SM}[k]^2 + \sqrt{\text{SM}[k]}^2}$$

Correction Factors

Code	Category	Explanation	Value	Error	Error(%)
0001	luminosity	CDF integrated luminosity	1990	50	2.6
0002	k-factor	cosmic_ph	0.81	0.05	6.1
0003	k-factor	cosmic_j	0.192	0.006	3.1
0004	k-factor	1γ1j photon+jet(s)	0.91	0.04	4.4
0005	k-factor	1γ2j	1.27	0.05	3.9
0006	k-factor	1γ3j	1.58	0.08	5.1
0007	k-factor	1γ4j+	1.99	0.16	8.1
0008	k-factor	2γ0j diphoton(+jets)	1.64	0.08	4.9
0009	k-factor	2γ1j	2.96	0.17	5.7
0010	k-factor	2γ2j+	1.2	0.09	7.5
0011	k-factor	W0j W (+jets)	1.37	0.03	2.3
0012	k-factor	W1j	1.32	0.03	2.3
0013	k-factor	W2j	2	0.05	2.5
0014	k-factor	W3j+	2.08	0.09	4.3
0015	k-factor	Z0j Z (+jets)	1.391	0.028	2.0
0016	k-factor	Z1j	1.23	0.04	3.2
0017	k-factor	Z2j+	1.02	0.04	3.9
0018	k-factor	2j $\hat{p}_T$<150 dijet	1.005	0.027	2.7
0019	k-factor	2j 150<$\hat{p}_T$	1.34	0.03	2.2
0020	k-factor	3j $\hat{p}_T$<150 multijet	0.945	0.025	2.6
0021	k-factor	3j 150<$\hat{p}_T$	1.48	0.04	2.7
0022	k-factor	4j $\hat{p}_T$<150	1.06	0.03	2.8
0023	k-factor	4j 150<$\hat{p}_T$	1.93	0.06	3.1
0024	k-factor	5j low	1.34	0.05	3.7
0025	k-factor	1b2j 150<$\hat{p}_T$	2.24	0.11	4.9
0026	k-factor	1b3j 150<$\hat{p}_T$	3.06	0.15	4.9
0027	misId	p(e→e) central	0.978	0.006	0.6
0028	misId	p(e→e) plug	0.965	0.007	0.7
0029	misId	p(μ→μ) CMUP+CMX	0.888	0.007	0.8
0030	misId	p(γ→γ) central	0.936	0.018	1.9
0031	misId	p(γ→γ) plug	0.86	0.016	1.9
0032	misId	p(b→b) central	0.971	0.021	2.2
0033	misId	p(γ→e) plug	0.06	0.003	5.0
0034	misId	p(q→e) central	7.07×10^{-5}	1.9×10^{-6}	2.7
0035	misId	p(q→e) plug	0.000785	1.2×10^{-5}	1.5
0036	misId	p(q→μ)	1.22×10^{-5}	6×10^{-7}	4.9
0037	misId	p(b→μ)	3.2×10^{-5}	1.1×10^{-5}	34.0
0038	misId	p(j→b) 25<p_T	0.0183	0.0002	1.1
0039	misId	p(q→τ)	0.0053	0.0001	1.9
0040	misId	p(q→γ) central	0.000269	1.4×10^{-5}	5.2
0041	misId	p(q→γ) plug	0.00048	6×10^{-5}	12.4
0042	trigger	p(e→trig) plug, p_T>25	0.838	0.007	0.8
0043	trigger	p(μ→trig) CMUP+CMX, p_T>25	0.92	0.004	0.4

Sleuth Algorithm

Sleuth variable:

$$\sum p_T \equiv \sum_i |\vec{p}_i| + \left|\overrightarrow{\text{uncl}}\right| + \left|\vec{\not{p}}\right|,$$

- Scan the sumPt spectrum in all final states and find the region with the most significant excess of data over SM.
- Perform pseudo-experiments to determine the probability that a statistical fluctuation of the background would yield an excess as significant as the one observed
- Takes into account the trials factor for looking at many places
- Discovery level significance set at 0.001 = 5σ effect

Method

Caution: This is NOT real data!



Scan all mass spectra with a window.

Window size follows mass resolution.

$$m = \sqrt{\left(\sum_i E_i\right)^2 - \left(\sum_i \vec{p}_i\right)^2} \Rightarrow \Delta m$$

- Consider bumps with ≥ 5 data events.
- Sidebands have to agree more than center, and not be too discrepant (5σ).

Statistical Significance





Each bump has a p-value

$$\sum_{n=d}^{\infty} \frac{b^n}{n!} e^{-b}$$

Most interesting bump: p-val$_{min}$

Use pseudo-data to find

P_a = The probability that a p-val ≤ p-val$_{min}$ would appear by coincidence.

$P_b = 1 - (1-P_a)^{\text{5000 distributions}}$

= probability at least one mass would have such a small P_a by coincidence.

Discovery threshold:

$P_b = 3\sigma \leftrightarrow P_a = 5\sigma$

Fast vs Slow method to estimate P_a



Expected P_a


Entries 2342
Mean 0.2204
RMS 1.453
Slow method
180
160
140
120
100
80
60
40
20
0
-10
-8
-6
-4
-2
0
2
4
6
8
10
σFast

The need for spike treatment



Spikes



Figure 4-10: *(Left)* The p-val of each bump candidate, as a function of the location of each window's center, along mass$(j1, j2)$ in final state $2j$ $\sum p_T < 400$ GeV. Bump candidates failing quality criteria have p-val=1. The most significant bump has p-val $\sim 10^{-6}$, which translates to $P_a \sim 3 \times 10^{-5}$ and $P_b \sim 0.15$, therefore all local excesses are insignificant. (Right) For demonstration, we apply the conservative anti-spike treatment to all bump candidates. The result of anti-spike treatment is to have larger p-values and the reduction of significance is greater in regions like around 400 GeV, where Monte Carlo statistics are poorer, therefore spikes contribute more.

Potential for Improvement

- Search for wider resonances
- Combine leptons & jet multiplicities
- Dynamic optimization of window width
- Use of only data

Jet -> Tau Fakes

Correction Factor p(q->tau) = 5.2E-3





jet -> photon fake

Correction Factor p(q->ph) central = 2.66E-4
Correction Factor p(q->ph) plug = 5.1E-4







jet -> photon fake





jet -> muon fake

Correction Factor p(q->mu) = 1.18E-5







Illustration of Fakes

jet -> ph fake

ph -> e fake

real photon